

Remarks on the Strong Coupling Constants in Heavy Hadron Chiral Lagrangians

Hai-Yang Cheng

Institute of Physics, Academia Sinica

Taipei, Taiwan 115, Republic of China

(January, 1997)

Abstract

In the heavy quark limit, there are three independent strong axial coupling constants in chiral Lagrangians for ground-state heavy hadrons: g in the meson sector, and g_1 , g_2 in the baryon sector. The coupling g_2 is extracted from recent CLEO measurements of $\Sigma_c^* \rightarrow \Lambda_c \pi$ to be $|g_2| = 0.57 \pm 0.10$, in agreement with the nonrelativistic quark model expectation but in 2σ deviation from the large- N_c argument. The parameter g_1 cannot be determined directly from strong decays of heavy baryons. Nevertheless, some information of g_1 can be learned from the radiative decay $\Xi_c'^{*0} \rightarrow \Xi_c^0 \gamma$, which is prohibited at tree level by SU(3) symmetry but induced by chiral loops. A measurement of $\Gamma(\Xi_c'^{*0} \rightarrow \Xi_c^0 \gamma)$ will yield two possible solutions for g_1 . Assuming the validity of the quark model relations among different coupling constants, the experimental value of g_2 implies that $|g| = 0.70 \pm 0.12$ and $|g_1| = 0.93 \pm 0.16$.

A most suitable framework for studying the low-energy dynamics of heavy hadrons is provided by the formalism in which heavy quark symmetry and chiral symmetry are synthesized [1, 2, 3]. In the limit of heavy quark symmetry, the heavy hadron chiral Lagrangian for strong interactions consists of three unknown strong coupling constants in the low-energy interactions between the Goldstone bosons and ground-state heavy hadrons: g in the meson sector, and g_1 , g_2 in the baryon sector. These three coupling constants are independent of the heavy quark species involved. In principle, the decays $D^* \rightarrow D\pi$, $\Sigma_c^* \rightarrow \Sigma_c\pi$ and $\Sigma_c \rightarrow \Lambda_c\pi$ can be utilized to determine the parameters g , g_1 and g_2 , respectively. Unfortunately, none of the decay rates of above decays has been measured. In particular, the decay $\Sigma_c^* \rightarrow \Sigma_c\pi$ (or $\Omega_c^* \rightarrow \Omega_c\pi$) is nowadays known to be kinematically prohibited since the mass difference between Σ_c^* and Σ_c is only of order 65 MeV. This means that the coupling constant g_1 cannot be determined directly from the strong decays of heavy baryons. Nevertheless, as pointed out in [1], the parameters g_1 and g_2 are related to g in the nonrelativistic quark model via ¹

$$g_1 = \frac{4}{3}g, \quad g_2 = -\sqrt{\frac{2}{3}}g. \quad (1)$$

In this short Letter we will extract g_2 from the recent CLEO measurement of $\Sigma_c^* \rightarrow \Lambda_c\pi$ decays. We then discuss its implications and the feasibility of measuring g_1 from the radiative decay $\Xi_c'^{*0} \rightarrow \Xi_c^0\gamma$.

The most general chiral-invariant low-energy interactions of ground-state heavy hadrons with the Goldstone bosons are given by (we follow the notation of [1])

$$\begin{aligned} \mathcal{L}_{\text{int}} = & f\sqrt{m_P m_{P^*}}(PA^\mu P_\mu^{*\dagger} + P_\mu^* A^\mu P^\dagger) + \frac{1}{2}g\epsilon_{\mu\nu\lambda\kappa}(P^{*\mu\nu}A^\lambda P^{*\kappa\dagger} + P^{*\kappa}A^\lambda P^{*\mu\nu\dagger}) \\ & + g_1\text{tr}(\bar{B}_6\gamma_\mu\gamma_5 A^\mu B_6) + g_2\text{tr}(\bar{B}_6\gamma_\mu\gamma_5 A^\mu B_{\bar{3}}) + \text{h.c.} \\ & + g_3\text{tr}(\bar{B}_{6\mu}^* A^\mu B_6) + \text{h.c.} + g_4\text{tr}(\bar{B}_6^{*\mu} A_\mu B_{\bar{3}}) + \text{h.c.} \\ & + g_5\text{tr}(\bar{B}_6^{*\nu}\gamma_\mu\gamma_5 A^\mu B_{6\nu}^*) + g_6\text{tr}(\bar{B}_{\bar{3}}\gamma_\mu\gamma_5 A^\mu B_{\bar{3}}), \end{aligned} \quad (2)$$

with $P_{\mu\nu}^{*\dagger} = D_\mu P_\nu^{*\dagger} - D_\nu P_\mu^{*\dagger}$, where P and P^* denote the ground-state 0^- and 1^- heavy mesons, respectively, $B_{\bar{3}}$ the spin- $\frac{1}{2}$ antitriplet baryon, B_6 the spin- $\frac{1}{2}$ sextet baryon, B_6^* the spin- $\frac{3}{2}$ baryon, and V_μ , A_μ are, respectively, the chiral vector and axial-vector fields. In the infinite heavy quark mass limit, heavy quark symmetry reduces the eight coupling constants to three independent ones, say g , g_1 and g_2 [1]:²

$$f = 2g, \quad g_3 = \frac{\sqrt{3}}{2}g_1, \quad g_5 = -\frac{3}{2}g_1, \quad g_4 = -\sqrt{3}g_2, \quad g_6 = 0. \quad (3)$$

¹The relation $g_1 = \frac{1}{3}g$ originally given in [1] is erroneous; the correction will appear in an erratum.

² In [4] the three independent parameters are denoted by g_1 , g_2 , g_3 , which are related to our notation by

$$g_1^{\text{Cho}} = g, \quad g_2^{\text{Cho}} = -\frac{3}{2}g_1, \quad g_3^{\text{Cho}} = -\sqrt{3}g_2.$$

Furthermore, there is only one independent coupling constant within the context of the non-relativistic quark model [see Eq. (1)].³ In the presence of $1/m_Q$ corrections, the heavy quark symmetry relations (3) are violated by the $1/m_Q$ heavy quark chromomagnetic operator $O_2 \sim \bar{Q}\sigma \cdot GQ$. Nevertheless, the O_2 -induced $1/m_Q$ corrections to the coupling constants still satisfy certain model-independent relations as elaborated in [6].

The theoretical estimate of the meson strong coupling constant g is rather diverse. Experimentally, the only constraint comes from the upper limit $\Gamma(D^{*+}) < 131$ keV set by ACCMOR Collaboration [7], which leads to $g < 0.74$. Theoretically, it is related in the nonrelativistic quark model to g_A^{ud} , the axial-vector coupling in the single quark transition $u \rightarrow d$, which has the value of 0.75 if the same model is required to produce the correct value of $g_A^N \simeq 1.25$ [1]. Relativistic effects of quark motion and quark spins are expected to reduce the value of g . Using the relativistic light-front quark model [8] we find $g \sim 0.50 - 0.55$, which is consistent with [9]. It has also been estimated in the approach of the QCD sum rule with the result [10]: $g \sim 0.21 - 0.39$. Using $g_A^{ud} = 0.75$, the coupling constants

$$g = 0.75, \quad g_1 = 1, \quad g_2 = -0.61 \quad (4)$$

can be regarded as the benchmarked values of g_i in the nonrelativistic quark model [1]. It should be stressed that, in spite of the experimental constraint $g < 0.74$, quark model calculations of the branching ratios of strong and radiative decays of charmed mesons based on $g = 0.75$ are consistent with experiment [11]. In the large- N_c approach, the baryon coupling constants are related to the nucleon axial coupling g_A^N via [12]

$$g_1 = g_A^N, \quad g_2 = -\frac{1}{\sqrt{2}} g_A^N. \quad (5)$$

The recent CLEO measurement of $\Sigma_c^* \rightarrow \Lambda_c \pi^\pm$ decays allows us to determine the coupling g_2 directly. From the baryon matrices

$$B_6 = \begin{pmatrix} \Sigma_c^{++} & \frac{1}{\sqrt{2}}\Sigma_c^+ & \frac{1}{\sqrt{2}}\Xi_c'^+ \\ \frac{1}{\sqrt{2}}\Sigma_c^+ & \Sigma_c^0 & \frac{1}{\sqrt{2}}\Xi_c'^0 \\ \frac{1}{\sqrt{2}}\Xi_c'^+ & \frac{1}{\sqrt{2}}\Xi_c'^0 & \Omega_c^0 \end{pmatrix}, \quad B_{\bar{3}} = \begin{pmatrix} 0 & \Lambda_c^+ & \Xi_c^+ \\ -\Lambda_c^+ & 0 & \Xi_c^0 \\ -\Xi_c^+ & -\Xi_c^0 & 0 \end{pmatrix}, \quad (6)$$

and B_6^* similar to B_6 , and

$$A_\mu = -\frac{1}{f_\pi} \partial_\mu \left(\frac{1}{2} \tau^a \pi^a \right) + \dots, \quad (7)$$

we obtain

$$A(\Sigma_c^* \rightarrow \Lambda_c \pi) = -i \frac{\sqrt{3}}{\sqrt{2}} \frac{g_2}{f_\pi} \bar{u}_{\Lambda_c} q_\mu u_{\Sigma_c^*}^\mu, \quad A(\Sigma_c \rightarrow \Lambda_c \pi) = -i \frac{\sqrt{3}}{\sqrt{2}} \frac{g_2}{f_\pi} \bar{u}_{\Lambda_c} q_\mu u_{\Sigma_c}, \quad (8)$$

³Using spin-flavor relativistic supermultiplet theory, it is shown in [5] that all the strong and electromagnetic interactions of ground-state hadrons can be expressed as a single coupling constant. However, the predicted strong decay widths of charmed baryons are in general quite different from ours. For example, the calculated rate $\Gamma(\Sigma_c \rightarrow \Lambda_c \pi) = 28$ keV in [5] is two orders of magnitude smaller than ours [see Eq. (17) below].

where u^μ is a Rarita-Schwinger vector-spinor for a spin- $\frac{3}{2}$ particle, q_μ is the pion 4-momentum, and $f_\pi = 93$ MeV. It is straightforward to show that

$$\begin{aligned}\Gamma(\Sigma_c^*(v) \rightarrow \Lambda_c(v)\pi) &= \frac{p_c^3}{6\pi} \left(\frac{\sqrt{3}g_2}{\sqrt{2}f_\pi} \right)^2 \frac{m_{\Lambda_c}}{m_{\Sigma_c^*}}, \\ \Gamma(\Sigma_c(v) \rightarrow \Lambda_c(v)\pi) &= \frac{p_c^3}{6\pi} \left(\frac{\sqrt{3}g_2}{\sqrt{2}f_\pi} \right)^2 \frac{m_{\Lambda_c}}{m_{\Sigma_c}},\end{aligned}\quad (9)$$

where p_c is the c.m. momentum of the final-state particle. In heavy quark effective theory, the velocity of Λ_c is taken to be the same as that of Σ_c^* or Σ_c . However, we will not apply heavy quark symmetry to the phase space due to large $1/m_Q$ corrections to p_c^3 ($m_{\Sigma_c} = m_{\Sigma_c^*}$ in heavy quark limit). If the realistic value of v_{Λ_c} is employed, the decay rates will become

$$\begin{aligned}\Gamma(\Sigma_c^*(v) \rightarrow \Lambda_c(v')\pi) &= \frac{p_c^3}{6\pi} \left(\frac{\sqrt{3}g_2}{\sqrt{2}f_\pi} \right)^2 \frac{(m_{\Sigma_c^*} + m_{\Lambda_c})^2 - m_\pi^2}{4m_{\Sigma_c^*}^2}, \\ \Gamma(\Sigma_c(v) \rightarrow \Lambda_c(v')\pi) &= \frac{p_c^3}{6\pi} \left(\frac{\sqrt{3}g_2}{\sqrt{2}f_\pi} \right)^2.\end{aligned}\quad (10)$$

It is easy to check that the zero-recoil relation $v \cdot v' = 1$ is a very good approximation for $\Sigma_c^* \rightarrow \Lambda_c \pi$ decay. As a consequence, one can apply either Eq. (9) or (10) to calculate $\Gamma(\Sigma_c^* \rightarrow \Lambda_c \pi)$. From the CLEO measurements [13]

$$\begin{aligned}\Gamma(\Sigma_c^{*++}) &= \Gamma(\Sigma_c^{*++} \rightarrow \Lambda_c^+ \pi^+) = 17.9_{-3.2}^{+3.8} \pm 4.0 \text{ MeV}, \\ \Gamma(\Sigma_c^{*0}) &= \Gamma(\Sigma_c^{*0} \rightarrow \Lambda_c^+ \pi^-) = 13.0_{-3.0}^{+3.7} \pm 4.0 \text{ MeV},\end{aligned}\quad (11)$$

and

$$\begin{aligned}m_{\Sigma_c^{*++}} &= m_{\Lambda_c} + (234.5 \pm 1.1 \pm 0.8) \text{ MeV}, \\ m_{\Sigma_c^{*0}} &= m_{\Lambda_c} + (232.6 \pm 1.0 \pm 0.8) \text{ MeV},\end{aligned}\quad (12)$$

we obtain ⁴

$$|g_2| = \begin{cases} 0.61 \pm 0.09, & \Sigma_c^{*++} \rightarrow \Lambda_c^+ \pi^+; \\ 0.53 \pm 0.11, & \Sigma_c^{*0} \rightarrow \Lambda_c^+ \pi^-. \end{cases}\quad (13)$$

We see that the average experimental value

$$|g_2| = 0.57 \pm 0.10 \quad (14)$$

is in agreement with the quark model prediction $|g_2| = 0.61$, but it deviates 2σ from the large- N_c argument: $|g_2| = 0.88$ [see Eq. (5)]. Note that our value of $|g_2|$ is slightly different from the result $|g_2| = (0.9 \pm 0.2)/\sqrt{3}$ obtained in [15] due mainly to the presence of the

⁴The same result for $|g_2|$ is also obtained by D. Pirjol and T.M. Yan [14].

kinematic factor ($m_{\Lambda_c}/m_{\Sigma_c^*}$) in Eq. (9). It is worth mentioning that $\Xi_c^{*\prime} \rightarrow \Xi_c \pi^\pm$ decays have also been seen by CLEO with the results [16]:

$$\begin{aligned}\Gamma(\Xi_c^{*\prime+}) &< 3.1 \text{ MeV}, & m_{\Xi_c^{*\prime+}} &= m_{\Xi_c^0} + (178.2 \pm 0.5 \pm 1.0) \text{ MeV}, \\ \Gamma(\Xi_c^{*\prime 0}) &< 5.5 \text{ MeV}, & m_{\Xi_c^{*\prime 0}} &= m_{\Xi_c^+} + (174.3 \pm 0.5 \pm 1.0) \text{ MeV}.\end{aligned}\quad (15)$$

We find $|g_2| < 0.64$ from the experimental limit on $\Gamma(\Xi_c^{*\prime+})$ and $|g_2| < 0.84$ from $\Gamma(\Xi_c^{*\prime 0})$. Applying the experimental value (14) leads to

$$\begin{aligned}\Gamma(\Xi_c^{*\prime+}) &= \Gamma(\Xi_c^{*\prime+} \rightarrow \Xi_c^+ \pi^0, \Xi_c^0 \pi^+) = (2.44 \pm 0.85) \text{ MeV}, \\ \Gamma(\Xi_c^{*\prime 0}) &= \Gamma(\Xi_c^{*\prime 0} \rightarrow \Xi_c^+ \pi^-, \Xi_c^0 \pi^0) = (2.51 \pm 0.88) \text{ MeV}.\end{aligned}\quad (16)$$

Note that we have neglected the effect of $\Xi_c - \Xi_c'$ mixing in calculations (for recent considerations, see [15, 17]). Therefore, the predicted total decay rate of $\Xi_c^{*\prime+}$ is very close to the current limit $\Gamma(\Xi_c^{*\prime+}) < 3.1 \text{ MeV}$ [16].

The decay width of Σ_c has not been measured. From Eqs. (9) and (14) we obtain

$$\Gamma(\Sigma_c^0 \rightarrow \Lambda_c^+ \pi^-) = (1.94 \pm 0.57) \text{ MeV}.\quad (17)$$

It is clear that the strong decay width of Σ_c is smaller than that of Σ_c^* by a factor of ~ 7 , although they will become the same in the limit of heavy quark symmetry. This is ascribed to the fact that the c.m. momentum of the pion is around 88 MeV in the decay $\Sigma_c \rightarrow \Lambda_c \pi$ while it is two times bigger in $\Sigma_c^* \rightarrow \Lambda_c \pi$.

Thus far we have neglected the electromagnetic contributions to the total decay width of heavy baryons. To check this, we note that the radiative decays are described by the amplitudes

$$\begin{aligned}A(B_6 \rightarrow B_3 + \gamma) &= i\eta_1 \bar{u}_3 \sigma_{\mu\nu} k^\mu \varepsilon^\nu u_6, \\ A(B_6^* \rightarrow B_3 + \gamma) &= i\eta_2 \epsilon_{\mu\nu\alpha\beta} \bar{u}_3 \gamma^\nu k^\alpha \varepsilon^\beta u^\mu, \\ A(B_6^* \rightarrow B_6 + \gamma) &= i\eta_3 \epsilon_{\mu\nu\alpha\beta} \bar{u}_6 \gamma^\nu k^\alpha \varepsilon^\beta u^\mu,\end{aligned}\quad (18)$$

where k_μ is the photon 4-momentum, ε_μ is the polarization 4-vector, and the coupling constants η_i can be calculated using the quark model [11]; some of them are

$$\begin{aligned}\eta_1(\Sigma_c^+ \rightarrow \Lambda_c^+) &= \frac{e}{6\sqrt{3}} \left(\frac{2}{M_u} + \frac{1}{M_d} \right), & \eta_2(\Sigma_c^{*+} \rightarrow \Lambda_c^+) &= \frac{e}{3\sqrt{6}} \left(\frac{2}{M_u} + \frac{1}{M_d} \right), \\ \eta_3(\Sigma_c^{*++} \rightarrow \Sigma_c^{++}) &= \frac{2\sqrt{2}e}{9} \left(\frac{1}{M_u} - \frac{1}{M_c} \right), & \eta_3(\Sigma_c^{*0} \rightarrow \Sigma_c^0) &= \frac{2\sqrt{2}e}{9} \left(-\frac{1}{2M_d} - \frac{1}{M_c} \right), \\ \eta_3(\Sigma_c^{*+} \rightarrow \Sigma_c^+) &= \frac{\sqrt{2}e}{9} \left(\frac{1}{M_u} - \frac{1}{2M_d} - \frac{2}{M_c} \right), & \eta_3(\Xi_c^{*\prime+} \rightarrow \Xi_c^+) &= \frac{e}{3\sqrt{6}} \left(\frac{2}{M_u} + \frac{1}{M_s} \right), \\ \eta_3(\Xi_c^{*\prime 0} \rightarrow \Xi_c^0) &= \frac{e}{3\sqrt{6}} \left(-\frac{1}{M_d} + \frac{1}{M_s} \right).\end{aligned}\quad (19)$$

Using the formulae ⁵

$$\begin{aligned}
\Gamma(B_6(v) \rightarrow B_{\bar{3}}(v) + \gamma) &= \eta_1^2 \frac{k^3}{\pi} \frac{m_f}{m_i}, \\
\Gamma(B_6^*(v) \rightarrow B_{\bar{3}}(v) + \gamma) &= \eta_2^2 \frac{k^3}{3\pi} \frac{m_f}{m_i}, \\
\Gamma(B_6^*(v) \rightarrow B_6(v) + \gamma) &= \eta_3^2 \frac{k^3}{3\pi} \frac{m_f}{m_i},
\end{aligned} \tag{20}$$

the light constituent quark masses

$$M_u = 338 \text{ MeV}, \quad M_d = 322 \text{ MeV}, \quad M_s = 510 \text{ MeV}, \tag{21}$$

from the Particle Data Group [18], and $M_c = 1.6 \text{ GeV}$, we obtain

$$\begin{aligned}
\Gamma(\Sigma_c^{*++} \rightarrow \Sigma_c^{++} \gamma) &= 1.4 \text{ keV}, & \Gamma(\Sigma_c^{*+} \rightarrow \Sigma_c^+ \gamma) &= 0.002 \text{ keV}, & \Gamma(\Sigma_c^{*0} \rightarrow \Sigma_c^0 \gamma) &= 1.2 \text{ keV}, \\
\Gamma(\Sigma_c^{*+} \rightarrow \Lambda_c^+ \gamma) &= 147 \text{ keV}, & \Gamma(\Sigma_c^+ \rightarrow \Lambda_c^+ \gamma) &= 88 \text{ keV}, \\
\Gamma(\Xi_c'^{*+} \rightarrow \Xi_c^+ \gamma) &= 54 \text{ keV}, & \Gamma(\Xi_c'^{*0} \rightarrow \Xi_c^0 \gamma) &= 1.1 \text{ keV}.
\end{aligned} \tag{22}$$

Hence, unlike the D^0 case where the radiative decay accounts for one third of the D^0 rate, the branching ratio of the radiative decays of charmed baryons is at most a few percent.

Assuming the validity of the quark model relations (1) for different coupling constants, it follows from (14) that

$$|g| = 0.70 \pm 0.12, \quad |g_1| = 0.93 \pm 0.16. \tag{23}$$

Since this value of g is substantially larger than the QCD sum rule estimate, it is of great importance to measure $\Gamma(D^*)$ to clarify the long-standing issue with g . As mentioned in passing, the parameter g_1 cannot be determined from the strong decays of heavy baryons. It has been advocated in [19] that a measurement of the branching ratio $\mathcal{B}(\Xi_c'^{*0} \rightarrow \Xi_c^0 \gamma)$ will determine $|g_1|$ directly. As pointed out in [20], the radiative decay $\Xi_c'^{*0} \rightarrow \Xi_c^0 \gamma$ is forbidden at tree level in SU(3) limit [see Eq. (19)]. In heavy baryon chiral perturbation theory, this radiative decay is induced via chiral loops where SU(3) symmetry is broken by the light current quark masses. A complete evaluation of leading chiral loops in [20] yields

⁵If the realistic value of v' is taken, Eq. (20) will become [11]

$$\begin{aligned}
\Gamma(B_6(v) \rightarrow B_{\bar{3}}(v') + \gamma) &= \eta_1^2 \frac{k^3}{\pi}, \\
\Gamma(B_6^*(v) \rightarrow B_{\bar{3}}(v') + \gamma) &= \eta_2^2 \frac{k^3}{3\pi} \frac{3m_i^2 + m_f^2}{4m_i^2}, \\
\Gamma(B_6^*(v) \rightarrow B_6(v') + \gamma) &= \eta_3^2 \frac{k^3}{3\pi} \frac{3m_i^2 + m_f^2}{4m_i^2}.
\end{aligned}$$

(see Eqs. (3.31), (3.39), (3.42) and (3.51) in [20])⁶

$$\begin{aligned} \eta_3(\Xi_c'^{*0} \rightarrow \Xi_c^0) = & -2\sqrt{\frac{3}{2}} \left[\frac{3}{8} g_1 g_2 a_1 (-\varepsilon_K + \varepsilon_\eta) + \frac{3}{2} g_2^2 a_2 (-\varepsilon_\pi + \varepsilon_K) \right. \\ & \left. + a_2 (-\varepsilon_\pi + \varepsilon_K) + \frac{3e}{128} \frac{g_1 g_2}{\pi f_\pi^2} (-m_\pi + m_K) \right], \end{aligned} \quad (24)$$

where

$$\varepsilon_P = \frac{1}{32\pi^2} \frac{m_P^2}{f_\pi^2} \ln \frac{\Lambda_\chi^2}{m_P^2}, \quad (25)$$

Λ_χ is a chiral-symmetry breaking scale of order 1 GeV, and a_1 , a_2 are coupling constants in the chiral Lagrangian for magnetic transitions of heavy baryons [11, 20]. It is clear that chiral-loop effects are nonanalytic in the forms of $m_q^{1/2}$ and $m_q \ln m_q$ (m_q being the current quark mass) and that a measurement of $\mathcal{B}(\Xi_c'^{*0} \rightarrow \Xi_c^0 \gamma)$ does not fix g_1 directly unless some model guidance on the parameters a_1 , a_2 is given. In the quark model a_i are simply related to the Dirac magnetic moments of light quarks. Writing (see Eq. (4.21) of [20])

$$a_1 = -\frac{e}{3}\beta, \quad a_2 = \frac{e}{2\sqrt{6}}\beta, \quad (26)$$

the parameter β is related in the nonrelativistic quark model to the constituent quark mass, $\beta = 1/M_q$ [11]. A study of radiative decays of the D mesons has confirmed this expectation [11, 21]. Using Eqs. (14), (24) and (26), we can utilize the measured $|\eta_3(\Xi_c'^{*0} \rightarrow \Xi_c^0)|$ to extract g_1 . By identifying (24) with the quark model prediction given by Eq. (19), we find two possible solutions: $|g_1| \approx 1.05$ and 0.31 . We see that one of the solutions for g_1 is in accord with the quark model value $g_1 = 1$. Of course, *a priori* there is no reason to expect that the two different approaches for radiative decays should agree with each other exactly since SU(3) violation is treated nonperturbatively in the quark model, while it is calculated in terms of a perturbative expansion in chiral effective Lagrangian theory. Nevertheless, the consistency between theory and model is very encouraging. It follows from (16) and (22) that the branching ratio of $\Xi_c'^{*0} \rightarrow \Xi_c^0 \gamma$ is of order 4×10^{-4} . Although it is probably difficult to measure this level of branching ratio in the near future, measurement of $\Gamma(\Xi_c'^{*0} \rightarrow \Xi_c^0 \gamma)$ may prove to be the only way to determine $|g_1|$, as accentuated in [19].

In summary, the strong coupling constant g_2 is extracted from the CLEO measurement of $\Sigma_c^* \rightarrow \Lambda_c \pi$ to be $|g_2| = 0.57 \pm 0.10$. In heavy baryon chiral perturbation theory, the radiative decay $\Xi_c'^{*0} \rightarrow \Xi_c^0 \gamma$ is induced by chiral loops where SU(3) symmetry is broken by the light current quark masses. A measurement of $\Gamma(\Xi_c'^{*0} \rightarrow \Xi_c^0 \gamma)$ will provide two possible solutions to g_1 . Assuming this radiative decay rate is the same as that estimated by the constituent

⁶ For simplicity we have neglected $\Xi_c - \Xi_c'$ mixing and Δm , the mass splitting between the sextet and antitriplet baryon multiplets, in loop calculations. The Δm correction to chiral effects of $m_q^{1/2}$ type is considered in [19], but it is not significant. Note that only one of the five chiral loop diagrams has been elaborated in [19] and contributions of order $m_q \ln m_q$ are not taken into account there.

quark model, we found that one of the solutions $|g_1| \approx 1.05$ is precisely what expected from the quark model. To conclude, the nonrelativistic quark model predictions for the strong coupling constants of ground-state heavy hadrons with Goldstone bosons: $g = 0.75$, $g_1 = 1$ and $g_2 = -0.61$, are consistent with all existing experiments.

ACKNOWLEDGMENTS

I am grateful to Tung-Mow Yan for discussions and for reading the manuscript. This work was supported in part by the National Science Council of ROC under Contract No. NSC86-2112-M-001-020.

References

- [1] T.M. Yan, H.Y. Cheng, C.Y. Cheung, G.L. Lin, Y.C. Lin, and H.L. Yu, *Phys. Rev.* **D46**, 1148 (1992); Erratum, to appear.
- [2] M.B. Wise, *Phys. Rev.* **D45**, 2188 (1992).
- [3] G. Burdman and J. Donoghue, *Phys. Lett.* **B280**, 287 (1992).
- [4] P. Cho, *Phys. Lett.* **B285**, 145 (1992).
- [5] R. Delbourgo and D. Liu, *Phys. Rev.* **D53**, 6576 (1996).
- [6] H.Y. Cheng, C.Y. Cheung, G.L. Lin, Y.C. Lin, T.M. Yan, and H.L. Yu, *Phys. Rev.* **D49**, 2490 (1994).
- [7] ACCMOR Collaboration, S. Barlary *et al.*, *Phys. Lett.* **B278**, 480 (1992).
- [8] H.Y. Cheng, C.Y. Cheung, and C.W. Hwang, IP-ASTP-04-96 [hep-ph/9607332], to appear in *Phys. Rev.* (1997).
- [9] P.J. O'Donnell and Q.P. Xu, *Phys. Lett.* **B336**, 113 (1994).
- [10] P. Colangelo, G. Nardulli, A. Deandrea, N. Di Bartolomeo, R. Gatto, and F. Feruglio, *Phys. Lett.* **B339**, 151 (1994); T.M. Aliev, D.A. Demir, E. Iltan, and N.K. Pak, *Phys. Lett.* **B351**, 339 (1995); V.M. Belyaev, V.M. Braun, A. Khodzhimirian, and R. Rückl, *Phys. Rev.* **D51**, 6177 (1995).

- [11] H.Y. Cheng, C.Y. Cheung, G.L. Lin, Y.C. Lin, T.M. Yan, and H.L. Yu, *Phys. Rev.* **D47**, 1030 (1993).
- [12] Z. Guralnik, M. Luke, and A.V. Manohar, *Nucl. Phys.* **B390**, 474 (1993); E. Jenkins, *Phys. Lett.* **B315**, 431 (1993).
- [13] CLEO Collaboration, G. Brandenburg *et al.*, CLNS 96/1427, CLEO 96-13 (1996).
- [14] T.M. Yan, private communication.
- [15] C.G. Boyd, M. Lu, and M.J. Savage, CMU-HEP96-15 [hep-ph/9612441].
- [16] CLEO Collaboration, P. Avery *et al.*, *Phys. Rev. Lett.* **75**, 4364 (1995); L. Gibbons *et al.*, *Phys. Rev. Lett.* **77**, 810 (1996).
- [17] T. Ito and Y. Matsui, *Prog. Theor. Phys.* **96**, 659 (1996).
- [18] Particle Data Group, *Phys. Rev.* **D54**, 1 (1996).
- [19] M. Liu, M.J. Savage, and J. Walden, *Phys. Lett.* **B369**, 337 (1996).
- [20] H.Y. Cheng, C.Y. Cheung, G.L. Lin, Y.C. Lin, T.M. Yan, and H.L. Yu, *Phys. Rev.* **D49**, 5857 (1994).
- [21] P. Cho and H. Georgi, *Phys. Lett.* **B296**, 408 (1992); J.F. Amundson, C.G. Boyd, E. Jenkins, M. Luke, A.V. Manohar, J.L. Rosner, M.J. Savage, and M.B. Wise *Phys. Lett.* **B296**, 415 (1992).